


Snake River Spring/Summer Chinook Captive Broodstock Rearing and Research

**Annual Report
2001**



DOE/BP-00004662-1

March 2003

This Document should be cited as follows:

Maynard, Desmond, W. McAuley, Thomas Flagg, "Snake River Spring/Summer Chinook Captive Broodstock Rearing and Research", Project No. 1996-06700, 24 electronic pages, (BPA Report DOE/BP-00004662-1)

Bonneville Power Administration
P.O. Box 3621
Portland, Oregon 97208

This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the author's and do not necessarily represent the views of BPA.

**SNAKE RIVER SPRING/SUMMER CHINOOK CAPTIVE
BROODSTOCK REARING AND RESEARCH, 2001**

Annual Report

Prepared by

Desmond J. Maynard
W. Carlin McAuley
and
Thomas A. Flagg

Resource Enhancement and Utilization Technologies Division
Northwest Fisheries Science Center
National Marine Fisheries Service
National Oceanic and Atmospheric Administration
Seattle, Washington

Prepared for

U.S. Department of Energy
Bonneville Power Administration
Environment, Fish and Wildlife
P.O. Box 3621
Portland, OR 97208-3621
Department of Energy

Project Number
Contract Number 96-AI-96441
Project 96-067-00

March 1, 2003

Executive Summary

In 1995, the National Marine Fisheries Service (NMFS), in cooperation with the Idaho Department of Fish and Game (IDFG), the Oregon Department of Fish and Wildlife (ODFW), and the Bonneville Power Administration (BPA) established captive broodstock programs to aid in the recovery of Snake River spring/summer chinook salmon (*Oncorhynchus tshawytscha*) listed as endangered under the U.S. Endangered Species Act (ESA). These programs were intended to provide safety nets for Salmon and Grande Ronde River Basins spring/summer chinook salmon stocks. They also provide a basis of examining the efficacy of captive rearing and captive breeding programs as tools for recovering listed salmonid populations. In years when no or few naturally produced fish return from the sea, captive fish and their progeny can be used to maintain populations in these two Snake River Basin tributaries.

The NMFS facility at Manchester, WA, provides the crucial seawater environment needed to culture anadromous salmonids during the marine phase of their life cycle. At the Manchester Research Station, the fish are cultured in 6.1m diameter circular tanks housed in a fully enclosed and secure building. The tanks are supplied with seawater that has been processed to eliminate most marine pathogens. The fish are fed a commercially prepared diet and held at densities and loading rates designed to maximize fish quality. When fish begin to mature, they are transferred to ODFW or IDFG freshwater facilities in Oregon and Idaho for final maturation. The states then release the mature fish (Idaho) or their progeny (Oregon) back into their native Snake River tributary waters in restoration efforts. In FY 2001, NMFS cultured 1996, 1997, 1998, and 1999 broodyear fish at its Manchester Facility. This report addresses program activities from September 1, 2000 to August 31, 2001.

TABLE OF CONTENTS

EXECUTIVE SUMMARY	ii
INTRODUCTION.....	1
FACILITIES.....	4
CAPTIVE BROODSTOCK FISH CULTURE PRACTICES	6
REINTRODUCTION STRATEGIES	12
REARING ACTIVITIES	13
SUMMARY	15
REFERENCES.....	16

INTRODUCTION

In spring 1995, the National Marine Fisheries Service (NMFS), US Fish and Wildlife Service (USFWS), Idaho Department of Fish and Game (IDFG), Oregon Department of Fish and Wildlife (ODFW), Confederated Tribes of the Umatilla Indian Reservation, and Nez Perce Tribe, initiated captive broodstocks as part of conservation efforts for ESA-listed stocks of Snake River spring/summer chinook salmon (*Oncorhynchus tshawytscha*). The Oregon captive broodstock program focuses on three stocks (upper Grande Ronde River, Catherine Creek, and Lostine River) captured as juveniles from the Grande Ronde River Basin. Idaho's Snake River captive broodstock program includes three stocks (Lemhi River, East Fork Salmon River, and West Fork Yankee Fork) captured as eyed eggs or juveniles from the Salmon River Basin. IDFG and ODFW requested that a portion of each broodstock be reared by NMFS in protective culture in seawater.

The captive broodstock concept differs from the conventional hatchery approach in that fish are maintained in captivity until they mature, rather than being released as conventional smolts that migrate to sea. "Captive breeding" fish are artificially spawned when they mature and their offspring are released back into their native waters as juveniles to supplement or restore the natural population. The relatively high egg-to-adult survival provided by the protective hatchery environment enables these programs to produce large numbers of juveniles in a single generation that can be used to jump start a depleted population (Flagg et al. 1995a,b). In addition, the relatively stable egg supply captive breeding programs provide helps ensure the success of supplementation efforts for depleted stocks. Oregon has chosen this traditional captive breeding approach where maturing fish are returned to Oregon, spawned, and the

offspring incubated, reared to the smolt stage, and then released into their native watersheds to restore their chinook salmon populations.

“Captive rearing”, the approach adopted by Idaho for their captive broodstocks differs from captive breeding, in that the mature adults are released back into their natal streams to spawn naturally. As with captive breeding, the high egg-to-smolt survival offered by the protective culture environment is used to rapidly increase the number of spawning adults above that which would occur if they had gone to sea. However, this approach is unproven and it is unclear whether hatchery-reared adults will retain the characteristics necessary for successful spawning in the wild. Because the success of achieving wild-fish attributes is uncertain, it is possible that in some years gametes may also be returned to Idaho to help maintain these gene pools using a captive breeding approach.

The NMFS Manchester Research Station (Manchester, WA) is one of the few West Coast facilities able to provide the crucial marine rearing needed to maintain the anadromous life history traits of Pacific salmon. Chinook salmon are an anadromous species that spawn in freshwater, spend one to two years rearing in streams and rivers, and then migrate to sea as smolts where they spend the majority (3-4 years) of their life (Healey 1991). When they begin to mature, adults migrate back to their natal rivers or streams to spawn. Salmon are thought to migrate to the sea, even though it imposes greater predation risk, because it provides better feeding and growing conditions. Thus, the potential survival reductions are offset by females producing more eggs and males developing enhanced sexual capabilities compared to population members that remain in freshwater. In addition, anadromous populations, with a portion of their members at sea each year, have an increased chance of survival when natural catastrophes (volcanic ash falls, drought years, etc.) eliminate all freshwater residents. Anadromy is the key

life history characteristic of Pacific salmon that enabled them to recolonize the vast vacant areas west of the Continental Divide after glacial recession. This evolutionary and ecological importance of anadromy behooves that conservation programs maintain captive Pacific salmon populations in seawater during the marine portion of their life cycle to eliminate possible selection against anadromous traits that might occur with full term freshwater rearing. The importance of seawater rearing during the marine phase of the salmon life cycle is recognized and called for by the NMFS Interim standards for the use of Captive Broodstocks (NMFS 1999).

The Snake River chinook salmon captive broodstocks are created by sourcing fish or eggs from natal river systems in Idaho and Oregon each year. Some of these fish are collected as parr in their first summer of life (ODFW AOP 2002), while others are obtained by pumping eyed eggs out of the gravel (Venditti et al. 2002). After collection, the fish or eyed eggs are transported to the IDFG Eagle Fish Hatchery or the ODFW Lookingglass Hatchery for freshwater rearing until they reach the smolt stage. Most of the smolts from the IDFG program and a third of the smolts from the ODFW program are transferred to the marine rearing facilities at Manchester. The fish are reared in seawater at Manchester until they begin to show signs of maturity. They are then transferred back to freshwater facilities in Idaho and Oregon for final maturation. The mature fish in the IDFG captive rearing program are released into enclosures in Salmon River streams to spawn naturally. ODFW artificially spawns the mature fish in its captive broodstock program at their Bonneville Hatchery to produce juveniles for smolt releases in their Grande Ronde Basin supplementation program. Investigators are comparing the relative success of the two reintroduction strategies and the effect of freshwater versus seawater rearing.

The use of captive broodstock fish to restore anadromous runs of Snake River chinook salmon is an action required to reach many of the objectives of the Northwest Power Planning

Council's (NPPC) Salmon Subbasin, and Grande Ronde Subbasin Summaries, and the 2000 NPPC Columbia Basin Fish and Wildlife Program. In addition, safety net captive broodstocks are among the Reasonable and Prudent Alternative (RPA 175, 176, 177) actions called for in the 2000 NMFS Biological Opinion, and are an ESA mandated item in the NMFS Recovery Plan for Snake River Salmon (Schmitten et al. 1995, 4.1a and 4.1b).

In August 1996, NMFS began a Bonneville Power Administration (BPA) funded project (Project 96-067-00) to rear Snake River spring/summer chinook salmon captive broodstocks in seawater at the NMFS Manchester Research Station. During 1997-2000, modifications were undertaken at Manchester to provide secure facilities for rearing of these ESA-listed fish. This included construction of a building housing a total of twenty 6.1-m diameter fiberglass rearing tanks, upgrade of the Manchester seawater pumping and filtration/sterilization systems to a total capacity of 4,542 L/min (1,200 gpm), and installation of an ozone depuration system.

Between 1996 and 2001, NMFS has captively-reared groups of Snake River spring/summer chinook salmon from Idaho and Oregon (Flagg et al. 1997, 1998; McAuley et al. 2000a,b). Maturity of these fish in captivity between fall 1996 and 2001 has resulted in more than 1,228 prespawning adults provided to IDFG and ODFW. In upcoming years, the cooperative broodstock program should continue to provide large numbers of animals for use in recovery efforts. Continuation of the cooperative captive broodstock programs may be critical to recovery of Snake River spring/summer chinook salmon.

FACILITIES

The marine rearing of the chinook captive broodstocks is conducted by NMFS at its Manchester Research Station near Manchester, Washington. The facility is located on nine

hectares of land adjoining Clam Bay in central Puget Sound. A major advantage of the site is the excellent seawater quality. Clam Bay is a major tidal mixing zone between Sinclair and Dyes Inlets to the west and waters of central Puget Sound to the east. Annual seawater temperature at the site normally ranges 7-13°C and salinity ranges 26-29 ppt. The high quality seawater environment, combined with a 250-m pier made available to the station by the EPA Region X Laboratory, make the Manchester Research Station an excellent site for salmon culture.

A 700-m-long pipeline from the end of the pier supplies about 4,542 L/min (1,200 gpm) of pumped seawater to the Station's land-based facilities. Water is pumped via two 50-hp vertical drive centrifugal pumps. The system is outfitted with back-up 50-hp pumps in case of primary pump failure. An alarm system monitors the pumps and electrical supply and is tied into an automatic dialer system linked to pagers and home and office telephones. Redundant emergency generators are automatically activated in the event of a power failure.

The marine rearing activities are conducted in Building 12 and 13. The 400-m² Building 12 contains six tan colored 4.1-m circular tanks inside the building and four gray colored 3.7-m fiberglass tanks in the adjoining fenced area outside the building. The 1,280-m² Building 13 houses 20 gray colored 6.1-m diameter circular fiberglass tanks. Tanks in both facilities are used for the project. In addition to these circular tanks, a number of portable circular tanks ranging in size from 0.8 to 2.3 m can be quickly set up for temporary fish holding.

The seawater supplied to these tanks is processed to eliminate pathogens. Filtering consists of primary sand filters containing number 20-grade sand that filters out all organic and inorganic material more than 20 microns in diameter. Water exiting the sand filters immediately enters a secondary cartridge filter system capable of filtering out all material more than 5 microns in diameter. The water then passes through UV-sterilizers for disinfection. Flow and

pressure sensors monitor flow through the seawater filtration/disinfection system.

The processed seawater is passed through packed column degassers to strip out any potential excess nitrogen and to boost dissolved oxygen levels. In addition, oxygen is continuously supplied to all rearing containers to both boost ambient dissolved oxygen levels to saturation and provide an emergency oxygen supply in the event of an unanticipated water flow interruption.

Prior to delivery to Building 13, water can be shunted through a chiller capable of reducing the temperature of a 1,135 L/min water supply by 5 degrees centigrade. Each rearing tank in Building 13 can be supplied with both ambient and chilled water through separate lines so that water temperature can be adjusted as needed. In most circumstances, water temperature will never be dropped more than 2 degrees centigrade below ambient. This will prevent an excessive temperature rise impacting fish health in the rare event of chiller failure.

The Manchester Research Station complies with Washington State quarantine certification standards by injecting ozone into all seawater exiting captive broodstock rearing areas. This enables fish from whirling disease-positive areas in the states of Idaho and Oregon to be moved in to the facility.

CAPTIVE BROODSTOCK FISH CULTURE PRACTICES

During the reporting period, fish were reared using standard fish culture practices and approved therapeutics (for an overview of standard methods see Leitritz and Lewis 1976, Piper et al. 1982, FRED 1983, Rinne et al. 1986, Shreck et al. 1995, and Pennell and Barton 1996). However, because captive broodstock husbandry for wild stocks is a new concept, the program generally used those practices that maximize fish quality and survival, rather than the number or

kg of fish produced as in traditional enhancement or commercial farming programs. All fish culture practices at the Manchester Research Station conformed to the husbandry requirements detailed in ESA Section 10 Propagation Permits for rearing of Idaho (Permit 972) and Oregon (Permit 973) stocks of ESA-listed Snake River spring/summer chinook salmon.

Both rearing and loading density were maintained on the low end of the scale to provide the best rearing environment possible. Generally, juvenile-to-adult rearing density in the tanks was maintained at under 8 kg/m³ (0.5 lbs/ft³) during most of the culture period, however, fish density ranged up to 15 kg/m³ (1.0 lbs/ft³) at maturity. Loading densities ranged from 0.29 kg/Lpm (2.5 lbs/gpm) to 0.84 kg/Lpm (7 lbs/gpm).

All tanks used for chinook captive broodstock rearing were completely covered with a taut 2.5 x 2.5 cm or smaller mesh nylon netting to prevent fish from jumping out. The energy absorbing properties of the nylon mesh minimizes injuries that can occur to fish when they leap against it. In addition to the mesh, half of each tank is covered with solid black fabric to provide a covered refuge area fish can move under when disturbed.

Fish were fed commercially prepared dry brood diets (Moore-Clark New Age Pacific and Pedigree Salmon Brood Special)¹. The daily ration ranged from 0.4 to 2.7% per day depending on estimated fish size and water temperature (Iwama 1996). Pellet size was determined from a chart provided by the manufacturer that is based on current guidelines for commercial aquaculture (Moore-Clark 1997). If necessary, pellet size was adjusted from the chart recommendation to ensure the smallest fish in the population could feed. Fish were fed by point

¹ Reference to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

source automatic feeders (Allen feeders or belt feeders). Each day, prior to loading the feeders, a small portion of the day's ration was broadcast over the surface to observe the fish's feeding response. The Allen feeders are programmed to disperse feed about 8 to 12 times a day during daylight hours. The feeding frequency was varied with both day length and fish size with smaller fish being fed 12 times per day and gradually reducing the frequency to 8 times per day as they grew. The belt feeders were only used for a short period with newly arriving smolts to disperse feed continuously over a 10-hour period.

A mild current (< 35 cm/sec) was generated in the rearing tanks by their circular shape, center drain, and a subsurface water jet inlet. This current primarily provided a self cleaning action and only a very slight exercise potential. At least once a week, bottom material that was not swept out of the tank by the current was removed by a combination of brushing and tank draining.

In May 2001, IDFG and ODFW transferred groups of 1999-brood Snake River spring/summer chinook salmon smolts to Manchester for rearing to adult (Table 1).

IDFG and ODFW both transferred the fish to Manchester as smolts in the spring of their second year of life. The first transfers from ODFW and IDFG were sentinels (10 fish from each stock) to determine the readiness of the population for seawater transition. When it was demonstrated these sentinels were surviving well and had begun to feed, the remainder of the fish were transferred. The main smolt groups were transitioned to seawater in 4-m diameter circular tanks in Building 12. The first step in this transitioning process was to add pathogen-free freshwater to each tank. The next step was to introduce the smolts to the tanks and then gradually add full strength Puget Sound seawater until all the freshwater had been displaced (an 8-12 h process). The postsmolts were generally held in these tanks through the summer before

being transferred to the 6.1-m diameter circular tanks in Building 13. Transfers of fish to Building 13 began when tanks became available as earlier year classes matured and were moved off-station. Similar transfer and rearing activities are planned for subsequent year-classes of fish.

Every effort was made to maintain the fish on low levels of natural lighting following the natural photoperiod. Both Building 12 and 13 had relatively small windows or translucent panels allowing a small percentage of ambient light to enter the building. In Building 13, additional lighting is provided by halogen lights that are automatically controlled to follow the natural photoperiod. These lights had a 30-minute ramp up (sunrise) and ramp down (sunset) feature. This ramping process reduces additional stress on the fish by avoiding the startle response associated with lights being instantaneously turned on at full intensity. In Building 12, additional lighting was provided by incandescent lights that are switched on and off as needed.

In general, the fish were handled with extreme care and kept in water to the maximum extent possible during transport and processing procedures. The transfer of ESA-listed fish was done with a dip net or fish transfer tube that held water during transfer to prevent the added stress of an out-of-water transfer. Additionally, the tank's center standpipe holds at least 15 cm of water depth in the tank when the external standpipe is pulled to minimize the chance of fish being accidentally dewatered during tank cleaning operations.

The Idaho stocks were not routinely sampled to minimize the impact of handling stress. Occasionally, the Idaho stocks were subsampled to assess fish size for determining proper feed size and ration. Sampling consisted of crowding the fish in a pool using two screens to form a wedge that confined the fish into a smaller area that facilitated netting. Netted fish were transferred to a tank containing an anaesthetic solution (MS222) and their PIT tag, weight, and length were recorded using the PITTag2 program (Pacific States Marine Fish Commission,

Gladstone, Oregon). The fish were then returned to the pool. After each pool was sampled, all equipment was disinfected with iodophor before sampling the next pool. The length (nearest 1 mm) and weight (nearest 0.1 g) of Oregon stocks was sampled four times a year by a team of NMFS, ODFW, and Nez Perce and Umatilla tribal personnel. ODFW is responsible for analyzing and reporting this sampling data.

The majority of inter-site transportation of juvenile or adult fish was be done by ODFW and IDFG, with assistance from NMFS. All transportation conducted by NMFS was done in a manner that emphasized fish health and safety. All transportation occurred in insulated containers and temperature was not allowed to rise more than 2 degrees C. The transport containers were supplied with a continuous oxygen supply that maintained dissolved oxygen at full saturation. The oxygen reservoir contained at least twice the oxygen needed to make the entire trip. The containers were loaded at no more than 0.06 kg/L (0.5 lbs/gallon). Additionally, drivers were equipped with cell phones, and had backup personnel ready to respond in event of equipment failure.

The captive broodstock fish were reared in seawater until they began to show signs of maturation. Fish culturists conducted maturation checks on fish in all groups suspected of having ripening adults. Maturation was determined by changes in skin sheen, skin coloration, and body morphology. However, few of those fish that have made the decision to mature in a given year can be identified before they would normally reenter freshwater. Therefore, in the spring of 2002, efforts will be made to determine if ultrasound scanning can be used to determine maturation status of fish earlier in the year. All fish determined to be maturing in seawater will be transferred to freshwater facilities in either Idaho or Oregon as soon as possible. In 2001, maturing adult fish from 1996- through 1999-year classes were returned to Idaho and Oregon for

use in recovery efforts for these stocks (Table 1). The transition from seawater back to freshwater is accomplished during transfer by transporting the maturing adults in a mixture of 1/4 seawater and 3/4 freshwater. This provides enough acclimation to allow the adults to be placed directly into freshwater tanks when reaching their respective freshwater captive broodstock rearing facilities.

Observable indices of fish health were checked daily by examining feeding response, external condition, and behavior of fish in each tank as initial indicators of developing problems. In particular, fish culturists observed for signs of lethargy, spiral swimming, side swimming, jumping, flashing, unusual respiratory activity, body surface abnormalities, and unusual coloration. A fish pathologist routinely monitored captive broodstock mortalities to determine cause of death. When a treatable pathogen was either detected or suspected in an Idaho stock, a veterinarian, in consultation with IDFG fish health staff, prescribed appropriate prophylactic and therapeutic drugs to control the problem. For the Oregon stocks, ODFW fish health staff prescribed appropriate treatments. Select mortalities were appropriately preserved for pathology, genetic, and other analyses. Specimens that were not vital to analysis were disposed of in a manner consistent with ESA permits.

NMFS provided daily staffing at the facility for protective culture of these fish, with constant electronic security and facilities monitoring. Staffing during some weekends and holidays was covered by drop-in site inspections. These inspections involved inspecting the facilities, loading feeders, removing and processing mortalities, and the mandatory daily backflushing of the filtration system. Electronic security and facilities monitoring were provided at all times. NMFS coordinated the details of rearing parameters for these fish with ODFW and IDFG through the CSCPTOC, and CCBPTOT process.

REINTRODUCTION STRATEGIES

The NMFS captive broodstock program has successfully generated prespawning adult chinook salmon to help restoration efforts for these ESA-listed endangered stocks.

Spring/summer chinook are being reared for the following two different reintroduction strategies as integral parts of the recovery process:

- Captive breeding (Oregon)—Adults are artificially spawned to provide offspring for smolt releases back into natal streams.
- Captive rearing (Idaho)—Adults are released into their natal streams for volitional spawning and natural production of offspring.

REARING ACTIVITIES

Brood-year 1996

In May 2001, remaining age-5 adults were checked for maturation status. A total of 31 prespawning Oregon adults were transferred to Bonneville Fish Hatchery (BFH) for spawning in early fall 2001. A total of seven age-5 Idaho adults were transferred to Eagle Fish Hatchery (EFH) for captive spawning, or release for volitional spawning in August 2001. Depending on stock, the percent survival in marine culture ranged from 0 to 80.6 for fish that matured by age 5 (Table 1).

Brood-year 1997

In May 2001, remaining age-4 adults were checked for maturation status. A total of 177 prespawning Oregon adults were transferred to BFH for spawning in early fall 2001. A total of 71 prespawning Idaho adults were transferred to EFH for captive spawning, or release for volitional spawning in August 2001. Depending on stock, the percent survival in marine culture ranged from 50.9 to 96.8 for fish that matured by age 4 (Table 1).

A preliminary investigation into the reproductive development rate of captive versus wild spring/summer chinook, conducted from May through August 2001 revealed varying amounts of abnormal ovary development in the captive stocks. Lethal samples were collected in May 2001 from 10 premature adults in the Lemhi River stocks, and five premature adults in the Catherine Creek stocks reared in seawater at Manchester, and 10 from the Catherine Creek stock reared in freshwater at BFH. In most cases, there was asymmetric development between the two ovaries, with atresia present in the seawater reared adults only (Penny Swanson, NMFS, NWFSC, Seattle, WA, 98112, Pers commun., May 2001). Reasons for this abnormal development are unknown, but distension of the stomach, due to retention of excessive amounts of seawater has

been suggested as a possible cause. This would not explain the asymmetry seen in the freshwater reared fish where the stomach distention was not present.

Brood-year 1998

In May 2001, remaining age-3 adults were checked for maturation status. A total of 138 prespawning Oregon adults were transferred to BFH for spawning in early fall 2001. A total of 87 prespawning Idaho adults were transferred to EFH for captive spawning, or release for volitional spawning in August 2001. Percent survival in marine culture for the various stocks of this year class ranged from 30.6 to 99.4 for fish that matured by age 3 (Table 1).

Brood-year 1999

In May 2001, ODFW transferred 332 smolts from Catherine Creek and Lostine River stocks to Manchester from Lookingglass Hatchery and IDFG transferred 630 smolts from Lemhi River, East Fork of the Salmon River, and safety-net stocks from the East Fork and West Fork Yankee Fork of the Salmon River to Manchester from EFH (Table 2). Sentinel fish from each stock were transferred approximately 7 days prior to ensure that the smolts were ready for seawater entry.

In July-August 2001, the Oregon stocks were checked for maturation status. A total of 27 prespawning age-2 precocial males were transferred to BFH for spawning in early fall 2001. Although the Idaho stocks were not checked for maturation, a total of 112 age-2 precocial males were present. Because representative groups were available at EFH for spawning or cryopreservation, the precocial males at Manchester were not utilized.

SUMMARY

The program has successfully maintained these ESA-listed populations in captivity and provided fish for use in restoration. The captive broodstock and captive rearing programs provide seawater rearing facilities that help ensure the retention of anadromous traits. This program, united with the IDFG and ODFW captive broodstocks should continue to provide for the restoration of self-sustaining natural runs of spring/summer chinook to the Salmon and Grande Ronde River Basins.

REFERENCES

- Flagg, T. A., C. V. W. Mahnken, and K. A. Johnson. 1995a. Captive broodstocks for recovery of depleted populations of Pacific salmon. *Am. Fish. Soc. Symp.* 15:81-90.
- Flagg, T. A., F. W. Waknitz, and C. V. W. Mahnken. 1995b. The captive broodstock concept: application to Pacific salmon. *In* T. A. Flagg and C. V. W. Mahnken (editors), *An assessment of the status of captive broodstock technology for pacific salmon*, pages 1-1 to 1-60. Report to Bonneville Power Administration, Contract DE-AI79-93BP55064.
- Flagg, T. A., M. R. Wastel, and W. C. McAuley. 1997. Manchester spring chinook broodstock project, progress report for 1996. Report to Bonneville Power Administration, Contract 96-BI-96441, 8 p.
- Flagg, T. A., M. R. Wastel, and W. C. McAuley. 1998. Manchester spring chinook broodstock project, progress report for 1997. Report to Bonneville Power Administration, Contract 96-BI-96441, 7 p.
- FRED. 1983. Fish culture manual. Alaska Department of Fish and Game, Division of Fisheries Rehabilitation, Enhancement and Development. Juneau, 90 p.
- Healey, M. C. 1991. Life history of chinook salmon (*Oncorhynchus tshawytscha*) *In* C. Groot and L. Margolis (editors), *Pacific Salmon life histories*, p 311-393. UBC Press, Vancouver.
- Iwama, G. K. 1996. Growth of salmonids. *In* W. Pennel and B. A. Barton (editors), *Principles of salmonid culture*, p. 467-516. Elsevier, Amsterdam.
- Leitritz, E., and R. C. Lewis. 1976. Trout and salmon culture (hatchery methods). *Calif. Dep. Fish Game Fish Bull.* 164, 197 p.
- McAuley, W. C., M. R. Wastel, and T. A. Flagg. 2000a. Manchester spring chinook broodstock project, progress report, 1998-1999. Report to Bonneville Power Administration, Contract 96-BI-96441, 8 p.
- McAuley, W. C., M. R. Wastel, and T. A. Flagg. 2000b. Manchester spring chinook broodstock project, progress report, 2000. Report to Bonneville Power Administration, Contract 96-AI-96441, 9 p.
- NMFS, 1999. Interim standards for the use of captive propagation technology in recovery of anadromous salmonids listed under the Endangered Species Act. NMFS Sustainable Fisheries Division, Portland, 18 p.
- ODFW, 2002. Grande Ronde Basin Captive Broodstock Status Update and Annual Operation Plan 2002. Oregon Department of Fish and Wildlife, 40p.

- Piper, R. G., I. B. McElwain, L. E. Orme, J. P. McCraren, L. G. Fowler, J. R. Leonard. 1982. Fish hatchery management. United States Department of Interior, Fish and Wildlife Service, Washington, D.C., 517 p.
- Pennell, W. and B. A. Barton. 1996. Principles of salmonid culture. Elsevier, Amsterdam, 1,070 p.
- Rinne, J. N., E. Johnson, B. L. Jensen, A. W. Ruger, and R. Sorenson. 1986. The role of hatcheries in the management and recovery of threatened and endangered fishes. *In* R. H. Stroud (editor), Fish culture in fisheries management, p. 271-285. Proceedings of a symposium on the role of fish culture in fisheries management. American Fisheries Society, Bethesda, Maryland.
- Schmittgen, R., W. Stelle, Jr., and R. P. Jones. 1995. Proposed Recovery Plan for Snake River Salmon, 347p., plus appendices.
- Schreck, C. B., M. S. Fitzpatrick, and K. P. Currens. 1995. Pacific salmon (*Oncorhynchus* spp.). *In* N. R. Bromage and R. J. Roberts (editors), Broodstock management and egg and larval quality, p. 197-219. Blackwell Science Ltd., Oxford.
- Venditti, D. A., C. Willard, C. Looney, P. Kline, and P. Hassemer. 2002. Captive rearing program for Salmon River chinook salmon. Annual progress report (2000) to the Bonneville Power Administration, Contract Number 00000167-00001.

Table 1. Inventory records for Oregon and Idaho stocks of ESA-listed 1994-, 1995-, 1996-, 1997-, 1998-, and 1999-brood spring/summer chinook salmon being reared from smolt to adult in seawater at the NMFS Manchester Research Station, September 2001.

Stock	Number transferred	Cumulative mortality	Precocial (%) ^b	Jacks (%) ^c	4, 5, and 6 year old adults (%) ^c	Number remaining	Survival ^a (%) ^d
1994-BROOD							
<u>IDAHO</u>							
Lemhi River	75	27	4 (5.4)	2 (2.9)	42 (95.5)	0	64.0
West Fork Yankee Fork	87	41	1 (1.2)	17 (23.9)	28 (75.7)	0	52.9
East Fork	75	26	4 (5.4)	17 (23.2)	28 (82.4)	0	65.3
<u>OREGON</u>							
Catherine Creek	167	32	15 (9.2)	38 (27.3)	82 (94.3)	0	80.8
Lostine River	160	70	5 (3.3)	20 (16.5)	65 (76.4)	0	56.3
1995-BROOD							
<u>IDAHO</u>							
Lemhi River	69	15	2 (3.4)	14 (24.6)	38 (62.3)	0	78.3
<u>OREGON</u>							
Catherine Creek	156	79	6 (5.0)	24 (23.3)	47 (65.3)	0	49.4
Lostine River	149	47	6 (4.2)	46 (35.3)	63 (79.4)	0	68.5
1996-BROOD							
<u>IDAHO</u>							
Lemhi River	110	64	0 (0.0)	12 (17.4)	34 (55.7)	0	41.8
West Fork Yankee Fork	60	50	8 (18.2)	0 (0.0)	2 (100.0)	0	16.7
East Fork	5	5	0 (0.0)	0 (0.0)	0 (0.0)	0	0.0
<u>OREGON</u>							
Catherine Creek	165	68	2 (1.3)	35 (23.0)	59 (52.2)	3	58.8
Lostine River	164	78	8 (5.2)	39 (26.0)	38 (55.1)	0	52.4
Grande Ronde	165	32	0 (0.0)	45 (30.2)	86 (93.5)	1	80.6

Table 1. Continued

Stock	Number Transferred	Cumulative mortality	Precocial (%) ^b	Jacks (%) ^c	4, 5, and 6 year old Adults (%) ^c	Number remaining	Survival ^a (%) ^d
1997-BROOD							
<u>IDAHO</u>							
Lemhi River	102	20	10 (10.2)	18 (20.9)	42 (76.4)	11	80.3
West Fork Yankee Fork	165	52	18 (12.0)	16 (14.5)	38 (46.3)	41	68.5
<u>OREGON</u>							
Catherine Creek	158	5	7 (4.5)	36 (24.0)	69 (62.2)	41	96.8
Lostine River	161	79	16 (10.0)	25 (18.5)	38 (77.6)	3	50.9
Grande Ronde	167	25	13 (7.8)	49 (32.7)	75 (84.3)	5	85.1
1998-BROOD							
<u>IDAHO</u>							
Lemhi River	158	23	12 (8.3)	25 (19.4)	--	98	85.4
West Fork Yankee Fork	193	34	25 (14.8)	35 (25.4)	--	99	82.4
East Fork	145	60	11 (12.8)	18 (24.0)	--	56	58.6
East Fork SN ^e	229	159	31 (15.3)	9 (22.0)	--	30	30.6
<u>OREGON</u>							
Catherine Creek	162	1	6 (3.7)	51 (32.7)	--	104	99.4
Lostine River	159	3	34 (21.7)	32 (26.2)	--	90	98.1
Grande Ronde	164	7	2 (1.2)	55 (34.8)	--	100	95.7
1999-BROOD							
<u>IDAHO</u>							
Lemhi River	210	25	47 (22.9)	--	--	138	88.1
West Fork Yankee Fork SN ^f	242	44	15 (6.5)	--	--	183	81.8
East Fork	113	3	44 (38.9)	--	--	66	97.3
East Fork SN ^e	65	9	6 (9.7)	--	--	50	86.2
<u>OREGON</u>							
Catherine Creek	166	3	8 (4.9)	--	--	155	98.2
Lostine River	166	1	19 (11.5)	--	--	142	99.4

^a Survival from seawater transfer.

^b Precocious fish from Oregon stocks were transferred to the ODFW Bonneville Fish Hatchery and milt cryopreserved. Precocious fish from Idaho stocks were removed from population and assayed for pathology, for all brood years except 1997. Precocial males from brood year 1997 were returned to the IDFG Eagle Hatchery and used in recovery evaluation.

^c Jacks, age 4, age 5, and age 6 adults from Oregon stocks were transferred to the ODFW Bonneville Fish Hatchery. Jacks, age 4, age 5, and age 6 adults from Idaho stocks were transferred to the IDFG Eagle Hatchery and used in recovery evaluations.

^d $100\% - ((\text{Number of mortalities}/\text{starting number}) * 100\%)$. Fish that survived to maturity are included as survivors.

^e East Fork Safety Net. These fish are second generation progeny of adults reared in captivity and spawned by IDFG at the Eagle Hatchery.

^f West Fork Yankee Fork Safety Net. These fish are second generation progeny of adults reared in captivity and spawned by IDFG at the Eagle Hatchery.

Table 2. Smolt transfer records for Oregon and Idaho stocks of ESA-listed 1999-brood spring/summer chinook salmon being reared from smolt to adult in seawater at the NMFS Manchester Research Station, September 2001.

Stock	Transfer Date	Number of Smolts	Average Weight (g)
IDAHO			
Lemhi River	5-10-01	210	48.1
West Fork Yankee Fork SN ^a	5-7-01	242	47.7
East Fork	5-7-01	113	50.7
East Fork SN ^b	5-10-01	65	35.6
OREGON			
Catherine Creek	5-15-01	166	21.2
Lostine River	5-15-01	166	20.5

^a West Fork Yankee Fork Safety Net. These fish are second generation progeny of adults reared in captivity and spawned by IDFG at the Eagle Hatchery.

^b East Fork Safety Net. These fish are second generation progeny of adults reared in captivity and spawned by IDFG at the Eagle Hatchery.